## FOR EMERGENCY USE ONLY FOR SALE AND USE IN SASKATCHEWAN UNTIL JULY 31, 2010. 0.4% STRYCHNINE BAIT

RESTRICTED

CONTAINS GRAIN THAT HAS BEEN MIXED WITH
2% LIQUID STRYCHNINE CONCENTRATE.
MAXIM CHEMICAL INTERNATIONAL LTD.
REGISTRATION NO. 28751 PEST CONTROL PRODUCTS ACT)
IN A RATIO OF

250 ML OF STRYCHNINE CONCENTRATE TO 1 KG OF GRAIN TO YIELD A GRAIN BAIT CONTAINING 0.4% STRYCHNINE.

TO BE USED ONLY IN THOSE RURAL MUNICIPALITIES IDENTIFIED BY SASKATCHEWAN AGRICULTURE AND FOOD AS HAVING A SEVERE INFESTATION OF RICHARDSON'S GROUND SQUIRRELS

READ THE LABEL AND ACCOMPANYING BROCHURE BEFORE USE.

DANGER POISON

CORROSIVE TO THE EYES

DO NOT USE FOR FOOD OR FEED.

THIS SEED HAS BEEN TREATED WITH STRYCHNINE.

RESTRICTED USE: For control of Richardson's ground squirrels (Spermophilus richardsonii).

DIRECTIONS FOR RESTRICTED USE:

TREATMENT OF BURROWS: 5-15 g (1-3 teaspoons) of strychnine treated grain must be placed deep in the burrow in such a manner that the bait is not visible from the surface. Following the bait application, the burrow entrance must be covered. Care must be taken to avoid spilling bait on soil surface. All spilled bait must be collected and disposed of by incineration or burial in a pit no less than 46 cm (approximately 18 inches) deep and then covered to prevent non-target poisonings. Toxic to all forms of life.

DO NOT SCATTER BAIT OUTSIDE BURROW.

NO ABOVE GROUND BAIT STATIONS MUST BE USED.